

The role of sea ice in atmosphere-ocean energy and momentum transfer *Insights from remote sensing*

Tom Armitage, Ron Kwok

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA

Georgy Manucharyan, Andrew F. Thompson

Environmental Science and Engineering, California Institute of Technology, Pasadena, CA

Alek Petty


Cryospheric Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD




Outline


- 1. Background
- 2. Surface stresses
- 3. Atmosphere-ice-ocean momentum transfer
- 4. Atmosphere-ice-ocean energy transfer
- 5. Eddy-ice interactions

- Satellite radar altimetry
 - Measures sea surface height over the global (wet) ocean




- Satellite radar altimetry
 - Measures sea surface height over the global (wet) ocean
 - Can estimate sea ice thickness








$$h_i = \left(\frac{\rho_w}{\rho_w - \rho_i}\right) h_f + \left(\frac{\rho_s}{\rho_w - \rho_i}\right) h_s$$






















- Arctic sea has cover has been changing rapidly
- What's been happening in the ocean?




Krishfield et al. (2014), JGR-Oceans

- Arctic sea has cover has been changing rapidly
- What's been happening in the ocean?
 - Beaufort Gyre freshwater content increase




Krishfield et al. (2014), JGR-Oceans



Giles et al. (2012), Nature Geoscience

- Arctic sea has cover has been changing rapidly
- What's been happening in the ocean?
 - Beaufort Gyre freshwater content increase
 - Increase in sea level



cm yr Giles et al. (2012), Nature Geoscience


- Arctic sea has cover has been changing rapidly
- What's been happening in the ocean?
 - Beaufort Gyre freshwater content increase
 - Increase in sea level
 - Speed up of surface currents

McPhee et al. (2013), J. Climate


2. Surface stresses

- Changes indicate surface stress increase/changes
 - Cannot be explained by atmospheric circulation change
- Very few observations of upper ocean to investigate this
- Currents often ignored in ice-ocean stress calculations

$$\vec{\tau}_{io} = \rho C_d (\vec{u}_i - \vec{u}_o) |\vec{u}_i - \vec{u}_o|$$



 Currents are similar magnitude to drift (sometimes faster!)



2. Surface stresses

 Can calculate sea ice-ocean stress and see the effect of non-zero upper ocean currents:


$$\vec{\tau}_{o} = (1 - C)\vec{\tau}_{ao} + C\vec{\tau}_{io}$$


$$\vec{\tau}_{ao} = \rho C_{da}\vec{u}_{a}|\vec{u}_{a}|$$

$$\vec{\tau}_{io} = \rho C_{do}(\vec{u}_{i} - \vec{u}_{o})|\vec{u}_{i} - \vec{u}_{o}|$$


• Ekman pumping (downwelling): $w_E = \frac{1}{
ho f} (
abla imes ec{ au}_o)$


Beaufort Gyre Ekman pumping with static ocean – nearly always negative







- Beaufort Gyre Ekman pumping with static ocean nearly always negative
- Including ocean current reduces Ekman pumping by nearly two thirds!
 - Always positive during the winter, means static ice, ocean drag induces upwelling






- Beaufort Gyre Ekman pumping with static ocean nearly always negative
- Including ocean current reduces Ekman pumping by nearly two thirds!
 - Always positive during the winter, means static ice, ocean drag induces upwelling






- Beaufort Gyre Ekman pumping with static ocean nearly always negative
- Including ocean current reduces Ekman pumping by nearly two thirds!
 - Always positive during the winter, means static ice, ocean drag induces upwelling




- Also important for the sea ice momentum budget
 - Significant difference in direction
 - Difference when including dynamic ocean is similar magnitude to absolute stress



4. Atmosphere–ice–ocean energy transfer

 Can also calculate work done on ocean geostrophic currents

$$W = \vec{\tau}_o \cdot \vec{u}_g$$




Wunsch (1998), J. Physical Oceanography

4. Atmosphere–ice–ocean energy transfer


 Can also calculate work done on ocean geostrophic currents


$$W = \vec{\tau}_o \cdot \vec{u}_g$$
 $W = (1 - C)\vec{\tau}_{ao} \cdot \vec{u}_g + C\vec{\tau}_{io} \cdot \vec{u}_g$
 W_{ao}
 W_{io}



4. Atmosphere-ice-ocean energy transfer


- Energy input dominated by direct from atmosphere
- Sea ice almost always acts to dissipate energy on average
- Major change in 2007 huge energy input and sustained increases after






4. Atmosphere-ice-ocean energy transfer

- Look at other terms in mechanical energy balance
- Find increased available potential energy
- Implies increased dissipation (by eddies) or advection out of the gyre






Wunsch (1998), J. Physical Oceanography

Sea ice momentum equation:


$$m\frac{\mathbf{D}\vec{u}_i}{\mathbf{D}t} = \vec{\tau}_{ai} + \vec{\tau}_{io} - mf(\vec{k} \times \vec{u}_i) - mg\nabla\eta + \nabla \cdot \sigma$$

- In MIZ conditions, summer conditions, m (i.e., thickness, concentration) gets small, ice (internal stress) gets weaker
- Get to the limit where ice-ocean stress, Coriolis, and internal stress are similar magnitude
- Find that ice motion is heavily influenced by mesoscale ocean eddies


- Animation shows high-res ice vorticity from ECCO
- Starts at end of winter
 - Thick, consolidated, strong ice
- Transitions into summer
 - Thinner, weaker, dispersed ice
- See clearly the impact of ocean eddies during summer




- Sentinel-1a/b provide ~daily coverage of the central Arctic Ocean
- Provide high-resolution (~10km) ice kinematics
- Eddy-ice motions clearly visible in MIZ
- Preliminary evidence of eddyice motions in summertime
 - Need to quantify uncertainty
 - Noisy in summer




170819_170820 Drift




Armitage et al., in prep



Shear




Armitage et al., in prep




25 20 15 (₁₋s w₀)


Vorticity



Armitage et al., in prep



Vorticity



Summary

- Accounting for upper ocean currents significantly changes ice-ocean stress estimates
- Ekman pumping is reduced ~two thirds when ocean currents included
 - Increased current act to modulate Ekman pumping, stabilize freshwater content
- Direct atmosphere-ocean work dominates energy input to the gyre;
 sea ice generally dissipates energy
 - Loss of sea ice mean more energy input, more eddies, more mixing
- Evidence for strong eddy-ice interactions in MIZ and during summer

Upper ocean currents are super important for ice-ocean interactions and need to be accounted for!

Data: Arctic Ocean: http://www.cpom.ucl.ac.uk/dynamic_topography/

Armitage et al. (2016), "Arctic sea surface height variability and change from satellite radar altimetry and GRACE, 2003-2014", JGR-Oceans, 121.

Armitage et al. (2017), "Arctic Ocean surface geostrophic circulation 2003-2014", The Cryosphere, 11.

Krishfield et al. (2014), "Deterioration of perennial sea ice in the Beaufort Gyre from 2003 to 2012 and its impact on the oceanic freshwater cycle", JGR-Oceans, 119.

Giles et al. (2012), "Western Arctic Ocean freshwater storage increased by wind-driven spin-up of the Beaufort Gyre", Nature Geoscience.

McPhee (2013), "Intensification of Geostrophic Currents in the Canada Basin, Arctic Ocean", Journal of Climate, 26.

Menghello et al. (2018), "Observations of seasonal upwelling and downwelling in the Beaufort Sea mediated by sea ice", Journal of Physical Oceanography, 48.

Dewey et al. (2018), "Arctic ice-ocean coupling and gyre equilibration observed with remote sensing", Geophysical Research Letters, 45.

Zhong et al. (2018), "Greater role of geostrophic currents in Ekman dynamics in the Western Arctic Ocean as a mechanism for Beaufort Gyre stabilization", *JGR-Oceans*, 123.



